

Date: August 17, 2018

From: J. A. Krieger

E-Mail: jonas.krieger@psi.ch

## MUSRFIT plug-in for simple $\beta$ -NMR resonance line shapes

This library contains useful functions to fit NMR and  $\beta$ -NMR line shapes. The functional form of the powder averages was taken from M. Mehring, Principles of High Resolution NMR in Solids (Springer 1983). The `libLineProfile` library currently contains the following functions:

### LineGauss

$$A(f) = e^{-\frac{4 \ln 2 (f-f_0)^2}{\sigma^2}} \quad (1)$$

Gaussian line shape around  $f_0$  with width  $\sigma$  and height 1.

MUSRFIT theory line: `userFcn libLineProfile LineGauss 1 2`

Parameters:  $f_0, \sigma$ .

### LineLaplace

$$A(f) = e^{-2 \ln 2 \left| \frac{f-f_0}{\sigma} \right|} \quad (2)$$

Laplaceian line shape around  $f_0$  with width  $\sigma$  and height 1.

MUSRFIT theory line: `userFcn libLineProfile LineLaplace 1 2`

Parameters:  $f_0, \sigma$ .

### LineLorentzian

$$A(f) = \frac{\sigma^2}{4(f-f_0)^2 + \sigma^2} \quad (3)$$

Lorentzian line shape around  $f_0$  with width  $\sigma$  and height 1.

MUSRFIT theory line: `userFcn libLineProfile LineLorentzian 1 2`

Parameters:  $f_0, \sigma$ .

### LineSkewLorentzian

$$A(f) = \frac{\sigma * \sigma_a}{4(f-f_0)^2 + \sigma_a^2}, \quad \sigma_a = \frac{2\sigma}{1 + e^{a(f-f_0)}} \quad (4)$$

Skewed Lorentzian line shape around  $f_0$  with width  $\sigma$ , height 1 and skewness parameter  $a$ .

MUSRFIT theory line: `userFcn libLineProfile LineSkewLorentzian 1 2 3`

Parameters:  $f_0, \sigma, a$ .

### LineSkewLorentzian2

$$A(f) = \begin{cases} \frac{\sigma_1^2}{4(f-f_0)^2 + \sigma_1^2}, & f < f_0 \\ \frac{\sigma_2^2}{4(f-f_0)^2 + \sigma_2^2}, & f > f_0 \end{cases} \quad (5)$$

Skewed Lorentzian line shape around  $f_0$  with height 1 and widths  $\sigma_1$ , and  $\sigma_2$ .

MUSRFIT theory line: `userFcn libLineProfile LineSkewLorentzian2 1 2 3`

Parameters:  $f_0, \sigma_1, \sigma_2$ .

### PowderLineAxialLor

$$A(f) = I_{ax}(f) \circledast \left( \frac{\sigma^2}{4f^2 + \sigma^2} \right) \quad (6)$$

Powder average of a axially symmetric interaction, convoluted with a Lorentzian.

$$I_{ax}(f) = \begin{cases} \frac{1}{2\sqrt{(f_{\parallel}-f_{\perp})(f-f_{\perp})}} & f \in (f_{\perp}, f_{\parallel}) \cup (f_{\parallel}, f_{\perp}) \\ 0 & \text{otherwise} \end{cases} \quad (7)$$

The maximal height of the curve is normalized to  $\sim 1$ .

MUSRFIT theory line: `userFcn libLineProfile PowderLineAxialLor 1 2 3`

Parameters:  $f_{\parallel}, f_{\perp}, \sigma$ .

### PowderLineAxialGss

$$A(f) = I_{ax}(f) \circledast \left( e^{-\frac{4 \ln 2 (f-f_0)^2}{\sigma^2}} \right) \quad (8)$$

Powder average of a axially symmetric interaction (Eq. 7), convoluted with a Gaussian. The maximal height of the curve is normalized to  $\sim 1$ .

MUSRFIT theory line: `userFcn libLineProfile PowderLineAxialGss 1 2 3`

Parameters:  $f_{\parallel}, f_{\perp}, \sigma$ .

### PowderLineAsymLor

$$A(f) = I(f) \circledast \left( \frac{\sigma^2}{4f^2 + \sigma^2} \right) \quad (9)$$

Powder average of a asymmetric interaction, convoluted with a Lorentzian. Assume without loss of generality that  $f_1 < f_2 < f_3$ , then

$$I(f) = \begin{cases} \frac{K(m)}{\pi \sqrt{(f-f_1)(f_3-f_2)}}, & f_3 \geq f > f_2 \\ \frac{K(m)}{\pi \sqrt{(f_3-f)(f_2-f_1)}}, & f_2 > f \geq f_1 \\ 0 & \text{otherwise} \end{cases} \quad (10)$$

$$m = \begin{cases} \frac{(f_2-f_1)(f_3-f)}{(f_3-f_2)(f-f_1)}, & f_3 \geq f > f_2 \\ \frac{(f-f_1)(f_3-f_2)}{(f_3-f)(f_2-f_1)}, & f_2 > f \geq f_1 \end{cases} \quad (11)$$

$$K(m) = \int_0^{\pi/2} \frac{d\varphi}{\sqrt{1 - m^2 \sin^2 \varphi}}, \quad (12)$$

where  $K(m)$  is the complete elliptic integral of the first kind. Note that  $f_1 < f_2 < f_3$  is not required by the code. The maximal height of the curve is normalized to  $\sim 1$ .

MUSRFIT theory line: `userFcn libLineProfile PowderLineAsymLor 1 2 3 4`

Parameters:  $f_1, f_2, f_3, \sigma$ .

### PowderLineAsymGss

$$A(f) = I(f) \circledast \left( e^{-\frac{4 \ln 2 (f-f_0)^2}{\sigma^2}} \right) \quad (13)$$

Powder average of a asymmetric interaction (Eq. 10-12), convoluted with a Gaussian. The maximal height of the curve is normalized to  $\sim 1$ .

MUSRFIT theory line: `userFcn libLineProfile PowderLineAsymGss 1 2 3 4`

Parameters:  $f_1, f_2, f_3, \sigma$ .